DirectGO™ PreMix-CE 2X PCR Master Mix

Operation Manual

Cat.: C108200-CE
Size: 1.25 mL (for 100 reactions)
Store at -20°C upon arrival
Research Use Only

I. Description:

2X DirectGO™ PreMix-CE is a premixed ready-to-use solution containing reaction buffer, dNTPs, glycerol, PCR enhancers, and Hot Start Taq DNA polymerase as a 2-fold concentration. Compared with conventional Taq polymerase, DirectGO™ Taq prevents non-specific amplification due to mis-priming or formation of primer dimers before thermal cycling, and it also possesses a 2X faster elongation rate than conventional Taq. The polymerase activity is restored during the initial denaturation step when amplification reactions are heated at 94-95°C for ten minutes. In addition, 2X DirectGO™ PreMix-CE is suitable for PCR amplification directly from whole blood, and samples from cell lines or tissues. You may skip any prior DNA purification steps, only applying a tiny amount of raw material directly into the PCR reaction, running it in a PCR machine, and loading it into an agarose gel for electrophoresis analysis. In addition, 2X DirectGO™ PreMix-CE is supplied with a specially optimized buffer that enables robust fluorescence signal for downstream capillary gel electrophoresis technology.

II. Applications:

1. Trace sample analysis
2. PCR - Karyotyping
3. Forensic examination
4. High - specificity PCR
5. High - throughput PCR

III. Storage Condition

-20℃	4℃	25℃	37℃
2 years	14 days	10 days	3 days

IV. Recommended PCR Condition:

Instruction on using whole blood as templates:

Heparin, EDTA, or citrate-treated whole blood samples are all suitable for DirectGO™ PreMix-CE. DNA amplification can be achieved in the presence of 1~10% blood.

Note:

For whole blood direct PCR, we recommend the reaction mix should be kept on ice before PCR amplification starts because a crude sample can result in false negative results from PCR.

Component	Volume per reaction (μL)
Raw material	X
10 μM forward Primer	0.5
10 μM reverse Primer	0.5
2X DirectGO™ PreMix-CE	12.5
Nuclease-Free Water	to 25
Total volume	25 μL

V. Thermocycling Conditions for a Routine PCR:

The recommended parameters may be optimized for each new primer-template pair for optimal specificity and amplification.

Cycles	Step	Temp.	Time
1	Initial denaturation	95℃	10 mins
25-35	Denaturation	95℃	30 sec
	Annealing	Primer Tm +3-5℃	30 sec
	Extension	72℃	15-30 sec/kb
1	Final Extension	72℃	5-10 mins
	Hold	14℃	--

VI. Experimental Data



Figure 1. Comparison of PCR amplification between DirectGO™ PreMix-CE and DirectGO™ PreMix after capillary electrophoresis analysis.

A and B lines were 5% and 10% blood samples in DirectGO™ PreMix-CE respectively; C and D lines were 5% and 10% blood samples in DirectGO™ PreMix; cartridge: S1 (BiOptic); sample injection: 4kv, 10sec; sample separation: 6kv, 300sec; alignment marker: 20bp-1kb (BiOptic).

Address: (23141) 4F, No.108-3,
Minquan Rd., Hsin-Tien District,
New Taipei City, Taiwan (R.O.C)

TEL: +886-2-2218-8726
FAX: +886-2-2218-8727
E-Mail: service@bioptic.com.tw